
Tullos Lumber Company, La Salle Parish, Louisiana, Zip Code 71371



The Louisiana Department of Health and Hospitals/Office of Public Health/Section of Environmental Epidemiology and Toxicology (SEET) cannot guarantee the accuracy of the information contained on this map and expressly disclaims liability for errors and omissions in its contents.

Louisiana Department of Environmental Quality Onsite Groundwater Sample Results, 1987-2005

Sample dates	Contaminant	N	Avg (mg/L) [‡]	Min (mg/L)	Max (mg/L)	N > MCL	% N > MCL
	[†] Arsenic (MCL* = 0.01 mg/L)	43	0.005	ND	0.048	6	14%
1992, 1995, 1997,	[†] Benzo(a)pyrene (MCL* = 0.0002 mg/L)	43	ND	ND	ND	0	0%
1999, 2000, 2001,	Chromium (MCL* = 0.1 mg/L)	43	0.026	ND	0.460	1	2%
2002, 2003	Copper (MCL** = 1.0 mg/L)	13	0.042	ND	0.350	0	0%
	Pentachlorophenol (MCL* = 0.001 mg/L)	0					

ND Not detected

- * Maximum Contaminant Level (National Primary Drinking Water Regulations, EPA)
- ** Secondary Maximum Contaminant Level (non-enforcable) (National Secondary Drinking Water Regulations, EPA)
- † Detection limits were greater than Maximum Contaminant Levels in some samples.
- ‡ A value of "0" was used to represent NDs when calculating the average (arithmetic mean).

Louisiana Department of Health and Hospitals, Office of Public Health, Safe Drinking Water Program Off-site (within 1/2 mile buffer zone) Drinking Water System Monitoring Results, 1991-2004 Aquifers: Williana-Bentley and Upland Terrace

Sample dates	Contaminant	N	Avg (mg/L) [‡]	Min (mg/L)	Max (mg/L)	N > MCL	% N > MCL
2000 2003 2004	[†] Arsenic (MCL* = 0.01 mg/L)	14	ND	ND	ND	0	0%
	Benzo(a)pyrene (MCL* = 0.0002 mg/L)	14	ND	ND	ND	0	0%
	Chromium (MCL* = 0.1 mg/L)	14	0.001	ND	0.010	0	0%
	Copper (MCL** = 1.0 mg/L)	2	3.60	2.70	4.50	2	100%
	Pentachlorophenol (MCL* = 0.001 mg/L)	14	ND	ND	ND	0	0%

ND Not detected

- ทิบ Not detected * Maximum Contaminant Level (National Primary Drinking Water Regulations, EPA)
- ** Secondary Maximum Contaminant Level (non-enforcable) (National Secondary Drinking Water Regulations, EPA)
- † Detection limits were greater than Maximum Contaminant Levels in some samples.
- ‡ A value of "0" was used to represent NDs when calculating the average (arithmetic mean).

Surveillance Epidemiology and End Results (SEER) Average Annual Cancer Incidence, 1999-2003 †

	La Salle	Parish		State of Louisiana				
Subgroup	ubgroup All Cancers Combined Lung* Bladder*		All Cancers Combined	Lung*	Bladder*			
White Males	558.2	112.2	42.3	595.5	108.7	39.1		
White Females	403.5	40.7	7.9	406.4	59.7	8.8		
Black Males	809.9	165.9	**	680.3	133.4	18.7		
Black Females	315.9	104.3	20.3	398.1	49.0	6.3		

- † November 2005 submission. Incidence rates (cases per 100,000 population per year) are age-adjusted to the 2000 US standard population.
- * Lung and bladder cancers are primarily associated with smoking. Lung cancer is also associated with exposure to asbestos, dioxin, metal fumes, radon, and certain chemicals. Bladder cancer is associated with analgesic abuse, chronic bladder problems, fluid intake, schistomiasis and chemicals such as azo dyes, benzidine.
- ** Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

2000 U.S. Census Bureau Zip Code Tabulation Area Demographic Information

Zip_Code	Total_Population	Male	Female	Under_Age_5	Over_64	White	Black	Other_Races	Ind_Below_Poverty_Level
71271	1 761	970 (9/ 40 4)	904 (9/ E0 G)	124 (9/7)	226 (0/ 42 4)	1 607	100	0	274 (0/44 7)

